

## **References:**

1. Barone JR, Schmidt WF. Effect of formic acid exposure on keratin fiber derived from poultry feather biomass. *Bioresource Technology* 2006; 97, 233- 242.
2. Cardamone JM. Investigating the microstructure of keratin extracted from wool: Peptide sequence (MALDI-TOF/TOF) and protein conformation (FTIR). *Journal of Molecular Structure* 2010; 969, 97-105.
3. Martínez-Hernández L, Velasco-Santos C, de Icaza M, Castaño V M. Grafting of methyl methacrylate onto natural keratin. *e-Polymers* 2003, no. 016.
4. Schmidt WF, Line, MJ. Physical and Chemical Structures of Poultry Feather Fiber Fraction in Fiber Process Development. In: *Proceedings of the TAPPI Nonwovens Conference 1996 March 11-13*; Charlotte, NC. Atlanta, GA: Technical Association of the Pulp and Paper Industry, 1996; pp.135–140.
5. Feughelman M. *Mechanical Properties and Structure of Alphakeratin Fibres*. University of New South Wales Press, Sydney 1997; 24–27.
6. Tesfaye T, Sithole B, Ramjugernath D. *Suitanable Chemistry and Pharmacy* 2018; 8: 38-49.
7. Nuutinen EM. *Feather characterization and processing*, Aalto University 2017.
8. Staroń P, Banach M, Kowalski Z. *Chemik* 2011; 65, 10: 1019-1026.