

## References

1. Yakhin OI, Lubyanov AA, Yakhin IA, Brown PH. Biostimulants in Plant Science: A Global Perspective. *Front. Plant Sci.* 2017; 7: 2049.
2. Bulgari R, Cocetta G, Trivellini A, Vernieri P, Ferrante A. Biostimulants and Crop Responses: A Review. *Biol Agric Hortic.* 2015; 31(1) 1–17.
3. Calvo P, Nelson L, Kloepper JW. Agricultural Uses of Plant Biostimulants. *Plant Soil.* 2014; 383: 3–41.
4. Toscano S, Romano D, Massa D, Bulgari R, Franzoni G, Ferrante A. Biostimulant Applications in Low Input Horticultural Cultivation Systems. *Italus Hortus.* 2018; 25(2): 27-36.
5. Roushphael Y, Colla G. Synergistic Biostimulatory Action: Designing the Next Generation of Plant Biostimulants for Sustainable Agriculture. *Front. Plant Sci.* 2018; 9:1655.
6. Cristiano G, Pallozzi E, Conversa G, Tufarelli V and De Lucia B. Effects of an Animal-Derived Biostimulant on the Growth and Physiological Parameters of Potted Snapdragon (*Antirrhinum Majus* L.). *Front. Plant Sci.* 2018; 9: 861.
7. Colla G, Roushphael Y, Canaguier R, Svecova E, Cardarelli M. Biostimulant Action of a Plant-Derived Protein Hydrolysate Produced Through Enzymatic Hydrolysis. *Front. Plant Sci.* 2014; 5: 448.
8. EBIC 2018. Available from: <http://www.biostimulants.eu/> (accessed 10.10.2019).
9. Wilson HT, Amirkhani M, Taylor AG. Evaluation of Gelatin as a Biostimulant Seed Treatment to Improve Plant Performance. *Front. Plant Sci.* 2018; 9:1006.
10. Kunicki E, Grabowska A, Sękara A, Wojciechowska R. The Effect of Cultivar Type, Time of Cultivation, and Biostimulant Treatment on the Yield of Spinach (*Spinacia oleracea* L.). *Folia Hortic.* 2010; 22:9–13.
11. Ziosi V, Zandoli R, Di Nardo A. Biological Activity of Different Botanical Extracts as Evaluated by Means of an Array of In Vitro and In Vivo Bioassays. *Acta Hortic.* 2013; 1009:61–66.
12. Parađiković N, Teklić T, Zeljković S, Lisjak M, Špoljarević M. Biostimulants Research in Some Horticultural Plant Species - A Review. *Food Energy Secur.* 2019; 8:e00162.
13. du Jardin P. Plant Biostimulants: Definition, Concept, Main Categories and Regulation. *Sci. Hortic.* 2015; 196: 3–14.
14. Drobek M, Frąc M, Cybulska J. Plant Biostimulants: Importance of the Quality and Yield of Horticultural Crops and the Improvement of Plant Tolerance to Abiotic Stress-A Review. *Agronomy.* 2019; 9: 335.
15. Nawrocka J, Małolepsza U. Diversity in Plant Systemic Resistance Induced by *Trichoderma*. *Biol Control.* 2013; 67: 149–156.
16. Rayirath P, Benkel B, Mark Hodges D, Allan-Wojtas P, MacKinnon S, Critchley AT, et al. Lipophilic Components of the Brown Seaweed, *Ascophyllum nodosum*, Enhance Freezing Tolerance in *Arabidopsis thaliana*. *Planta.* 2009; 230(1): 135–47.
17. Bradáčová K, Weber NF, Morad-Talab N, Asim M, Imran M, Weinmann M, et al. Micronutrients (Zn/Mn), Seaweed Extracts, and Plant Growth-Promoting Bacteria as Cold-Stress Protectants in Maize. *Chem Biol Technol Agric.* 2016; 3(1): 19.

18. Elansary HO, Skalicka-Woźniak K, King IW. Enhancing Stress Growth Traits as Well as Phytochemical and Antioxidant Contents of Spiraea and Pittosporum Under Seaweed Extract Treatments. *Plant Physiol Biochem.* 2016; 105: 310–20.
19. Zhang X, Ervin EH. Cytokinin-Containing Seaweed and Humic Acid Extracts Associated with Creeping Bentgrass Leaf Cytokinins and Drought Resistance. *Crop Sci.* 2004; 44(5): 1737.
20. García AC, Santos LA, Izquierdo FG, Sperandio MVL, Castro RN, Berbara RLL. Vermicompost Humic Acids as an Ecological Pathway to Protect Rice Plant Against Oxidative Stress. *Ecol Eng.* 2012; 47: 203–8.
21. El-Nemr MA, El-Desuki M, El-Bassiony AM, Fawzy ZF. Response of Growth and Yield of Cucumber Plants (*Cucumis Sativus L.*) To Different Foliar Applications of Humic Acid and Bio-Stimulators. *Aust J Basic Appl Sci.* 2012; 6:630–637.
22. Anjum SA, Wang L, Farooq M, Xue L, Ali S. Fulvic Acid Application Improves the Maize Performance Under Well- Watered And Drought Conditions. *J Agron Crop Sci.* 2011b; 197: 409–417.
23. Peng A, Xu Y, Wang ZJ. The Effect of Fulvic Acid on the Dose Effect of Selenite on the Growth of Wheat. *Biol Trace Elem Res.* 200183: 275–279.
24. Rady MM, Abd El-Mageed TA, Abdurrahman HA, Mahdi AH. Humic Acid Application Improves Field Performance of Cotton (*Gossypium Barbadense L.*) Under Saline Conditions. *J. Anim. Plant Sci.* 2016, 26(2): 487-493.
25. Belopukhov SL, Grishina EA, Dmitrevskaya II, Lukomets VM, Uschapovsky IV. Effect of Humic-Fulvic Complex on Flax Fiber and Seed Yield Characteristics. *Известия ТГУА.* 2015; 4: 631.811.98.
26. Colla G, Rouphael Y, Di Mattia E, El-Nakheel C, Cardarelli M. Co-Inoculation of *Glomus Intraradices* And *Trichoderma Atroviride* Acts as a Biostimulant to Promote Growth, Yield and Nutrient Uptake of Vegetable Crops. *J. Sci. Food Agric.* 2015a; 95: 1706-1715.
27. Azarmi R, Hajieghrari B, Giglou A. Effect of Trichoderma Isolates on Tomato Seedling Growth Response and Nutrient Uptake. *Afr. J. Biotechnol.* 2011; 10(31): 5850-5855.
28. Nawrocka J, Małolepsza U, Szymczak K, Szczech M. Involvement of Metabolic Components, Volatile Compounds, PR Proteins, and Mechanical Strengthening in Multilayer Protection of Cucumber Plants Against *Rhizoctonia Solani* Activated by *Trichoderma Atroviride* TRS25. *Protoplasma.* 2018; 255, 359–373.
29. Ertani A, Schiavon M, Muscolo A, Nardi S. Alfalfa Plant-Derived Biostimulant Stimulate Short-Term Growth of Salt Stressed *Zea Mays L.* Plants. *Plant Soil.* 2013; 364, 145–158.
30. Zhu K, Zhou H, Qian H. Antioxidant and Free Radical-Scavenging Activities of Wheat Germ Protein Hydrolysates (WGPH) Prepared with Alcalase. *Process Biochem.* 2006; 41(6): 1296–302.
31. Lucini L, Rouphael Y, Cardarelli M, Canaguier R, Kumar P, Colla G. The Effect of a Plant-Derived Biostimulant on Metabolic Profiling and Crop Performance of Lettuce Grown Under Saline Conditions. *Sci Hortic.* 2015; 23(182): 124–33.
32. Lachhab N, Sanzani SM, Adrian M, Chilitz A, Balacey S, Boselli M. et al. Soybean and Casein Hydrolysates Induce Grapevine Immune Responses and Resistant Against *Plasmopara Viticola*. *Front. Plant Sci.* 2014; 5: 716.

33. Craigie JS. Seaweed Extract Stimuli in Plant Science and Agriculture. *J Appl Phycol.* 2011; 23: 371–393.
34. Jansa J, Gryndler M. Biotic Environment of the Arbuscular Mycorrhizal Fungi in Soil, In: Koltai H, Kapulnik Y, editors. *Arbuscular Mycorrhizas: Physiology and Function* Springer Science+Business Media B.V. 2010; pp. 223.
35. Khan W, Rayirath UP, Subramanian S. et al. Seaweed Extracts as Biostimulants of Plant Growth and Development. *J Plant Growth Regul.* 2009; 28:386–399.
36. Zanin L, Tomasi N, Cesco S, Varanini Z and Pinton R. Humic Substances Contribute to Plant IronNutrition Acting as Chelators and Biostimulants. *Front. Plant Sci.* 2019; 10: 675.
37. Tipping E. Cation Binding by Humic Substances. Cambridge: Cambridge University Press. 2002; 1–434.
38. Trevisan S, Franciosi O, Quaggiotti S, Nardi S. Humic Substances Biological Activityat the Plant-Soil Interface. From Environmental Aspects to Molecular Factors. *Plant Signal Behav.* 2010; 5:6, 635-643.
39. Szczech M, Nawrocka J, Felczyński K, Małolepsza U, Sobolewski J, Kowalska B, Maciorowski R, Jas K, Kancelista A. *Trichoderma Atroviride* TRS25 Isolate Reduces Downy Mildew and Induces Systemic Defence Responses in Cucumber in Field Conditions. *Sci. Hortic.* 2017; 224: 17-26.
40. Karolev N, Rav DD, Elad Y. The Role of Phytohormones in Basal Resistance and *Trichoderma*-Induced Systemic Resistance to *Botrytis Cinerea* in *Arabidopsis Thaliana*. *Biol Control.* 2008; 53: 667-683.
41. Djonović S, Pozo MJ, Dangott LJ, Howell CR, Kenerley CM. Sm1, A Proteinaceous Elicitor Secreted by The Biocontrol Fungus *Trichoderma Virens* Induces Plant Defense Responses and Systemic Resistance. *Mol Plant Microbe Interact.* 2006; 7: 838-853.
42. Martínez-Medina A, Fernández I, Sánchez-Guzmán M, Jung SC, Pascual JA, Pozo MJ. Deciphering the Hormonal Signalling Network Behind the Systemic Resistance Induced by *Trichoderma Harzianum* in Tomato. *Front. Plant Sci.* 2013; 4(206): 1-12.
43. Nagaraju A, Sudisha J, Murthy SM. Ito SI. Seed Priming with *Trichoderma Harzianum* Isolates Enhances Plant Growth and Induces Resistance Against *Plasmopara Halstedii*, an Incitant of Sunflower Downy Mildew Disease. *Australas Plant Path.* 2012; 41: 609-620.
44. De Palma M, D'Agostino N, Proietti S, Bertini L, Lorito M, Ruocco M, Caruso C, Chiusano ML, Tucci M. Suppression Subtractive Hybridization Analysis Provides New Insights into the Tomato (*Solanum Lycopersicum L.*) Response to the Plant Probiotic Microorganism *Trichoderma Longibrachiatum* MK1. *J. Plant Physiol.* 2016; 190: 79-94.
45. Martinez C, Blanc F, Le Claire E, Besnard O, Nicole M, Baccou JC. Salicylic acid and Ethylene Pathways are Differentially Activated in Melon Cotyledons by Active or Heat-Denatured Cellulase from *Trichoderma Longibrachiatum*. *Plant Physiology.* 2001; 127: 334-344.
46. [Segarra G](#), [Casanova E](#), [Avilés M](#), [Trillas I](#). *Trichoderma Asperellum* Strain T34 Controls *Fusarium* Wilt Disease in Tomato Plants in Soilless Culture Through Competition for Iron. *Microb Ecol.* 2010; 59(1): 141-9.

47. Harman GE, Herrera-Estrella AH, Horwitz BA, Lorito M. Special Issue: *Trichoderma*— from Basic Biology to Biotechnology. *Microbiology*. 2012; 158: 1-2.
48. Hermosa R, Viterbo A, Chet I, Monte E. Plant-Beneficial Effects of *Trichoderma* and of its Genes. *Microbiology*. 2012; 158: 17-25.
49. Sharma S, Kour D, Rana KL, Dhiman A, Thakur S, Thakur P, Thakur S, Thakur N, Sudheer S, Yadav N, Yadav AN, Rastegari AA, Singh K. *Trichoderma*: Biodiversity, Ecological Significances, and Industrial Applications. In: Yadav AN, Mishra S, Singh S, Gupta A. editors. Recent Advancement in White Biotechnology Through Fungi. *Fungal Biology*. 2019; pp. 85-109.
50. Błaszczyk L., Siwulski M., Sobieralski K., Lisiecka J., Jędryczk M. *Trichodermaspp.* – Application and Prospects for use in Organic Farming and Industry. *J. Plant Prot. Res.* 2014; 54(4): 309-317.
51. Hýsek J, Vach M, Brožová J, Sychrová E, Civínová M, Nedělník J, Hrubý J. The Influence of the Application of Mineral Fertilizers with the Biopreparation Supresivit (*Trichoderma Harzianum*) on the Health and the Yield of Different Crops. *Arch Phytopathology Plant Protect.* 2002; 35(2): 115-124.
52. López-Bucio J, Pelagio-Flores R, Herrera-Estrella A. Trichoderma as Biostimulant: Exploiting the Multilevel Properties of a Plant Beneficial Fungus. *Sci Hortic.* 2015; 30(196): 109–23.
53. Schiavon M, Pizzeghello D, Muscolo A, Vaccaro S, Francioso O, Nardi S. High Molecular Size Humic Substances Enhance Phenylpropanoid Metabolism in Maize (*Zea Mays L.*). *J. Chem. Ecol.* 2010; 36, 662–669.
54. Vranova V, Rejsek K, Skene KR, Formanek P. Non-Protein Amino Acids: Plant, Soil and Acosystem Interactions. *Plant Soil.* 2011; 342:31–48.
55. Tegeder M. Transporters for Amino Acids in Plant Cells: Some Functions and Many Unknowns. *Curr. Opin. Plant. Biol.* 2012; 15: 315–321.
56. Kauffman GL, Kneival DP, Watschke TL. Effects of Biostimulant on the Heat Tolerance Associated with Photosynthetic Capacity, Membrane Thermostability, and Polphenol Production Ofperennial Ryegrass. *CropSci.* 2007; 47:261–267.
57. Sharma A, Shankhdhar D, Shankhdhar SC. Enhancing Grain Iron Content Of Rice By The Application Of Plant Growth Promoting Rhizobacteria. *Plant Soil Environ.* 2013; 59: 89–94.
58. Norrie J, Keathley JP. Benefits of *Ascophyllum Nodosum* Marine-Plant Extract Applications to “Thompson Seedless” Grape Production. *Acta Hortic.* 2006; 727:243–247.
59. Hamza B, Suggars A. Biostimulants: Myths and Realities. *Turfgrass Trends.* 2001; 10:6–10.
60. Canellas LP, Olivares FL, Aguiar NO, Jones DL, Nebbioso A, Mazzei P, et al. Humic and Fulvic Acids as Biostimulants in Horticulture. *Sci Hortic.* 2015; 30(196):15–27.
61. Baldotto MA, Baldotto LEB. Gladiolus Development in Response to Bulb Treatment with Different Concentrations of Humic Acids. *Rev Ceres.* 2013; 60:138–142.
62. Canellas LP, Olivares FL, Okorokova-Façanha AL, Façanha AR. Humic Acids Isolated from Earthworm Compost Enhance Root Elongation, Lateral Root

- Emergence, and Plasma Membrane HC-Atpase Activity In Maize Roots. *Plant Physiol.* 2002; 130: 1951–1957.
- 63. Nardi S, Pizzeghello D, Muscolo A, Vianello A. Physiological effects of Humic Substances on Higher Plants. *Soil Biol. Biochem.* 2002; 34, 1527–1536.
  - 64. García AC, Olaetxea M, Santos LA, Mora V, Baigorri R, Fuentes M. et al. Involvement of Hormone-and ROS-Signaling Pathways in the Beneficial Action of Humic Substances on Plants Growing Under Normal and Stressing Conditions. *BioMed. Res. Int.* 2016b; 2016: 3747501.
  - 65. García AC, Santos LA, de Souza LGA, Tavares OCH, Zonta E, Gomes ETM. et al. Vermicompost Humic Acids Modulate the Accumulation and Metabolism of ROS in Rice Plants. *J. Plant Physiol.* 2016c; 192, 56–63.
  - 66. Ramos AC, Dobbss LB, Santos LA, Fernandes MS, Olivares FL, Aguiar NO et al. Humic Matter Elicits Proton and Calcium Fluxes and Signalling Dependent on  $\text{Ca}^{2+}$ -Dependent Protein Kinase (CDPK) at Early Stages of Lateral Plant Root Development. *Chem. Biol. Tech. Agr.* 2015; 2: 3.
  - 67. Mora V, Baigorri R, Bacaicoa E, Zamarreño AM, García-Mina JM. The Humic Acid-Induced Changes in the Root Concentration of Nitric Oxide, IAA and Ethylene do not Explain the Changes in Root Architecture Caused by Humic Acid in Cucumber. *Environ. Exp. Bot.* 2012; 76, 24–32.
  - 68. Zandonadi DB, Santos MP, Dobbss LB, Olivares FL, Canellas LP, Binzel ML. et al. Nitric Oxide Mediates Humic Acids Induced Root Development and Plasma Membrane H+-Atpase Activation. *Planta*. 2010; 231, 1025–1036.
  - 69. Baldotto LEB, Baldotto MA, Giro VB, Canellas LP, Olivares FL, Bressan-Smith R. Performance of Pineapple ‘Vito’Ria’ in Response to the Application of Humic Acids During Acclimatization. *Rev Bras Ciéncia Solo.* 2009; 33:979–990.
  - 70. Chen Y, Clapp CE, Magen H. Mechanisms of Plant Growth Stimulation by Humic Substances: The Role of Organo-Iron Complexes. *Soil Sci Plant Nutr.* 2004; 50: 1089–1095.
  - 71. Harman GE, Howell CR, Viterbo A, Chet I, Lorito M. *Trichoderma* Species – Opportunistic, Avirulent Plant Symbionts. *Nat. Rev. Microbiol.* 2004; 2(1): 43–56.
  - 72. Contreras-Cornejo HA, Macías-Rodríguez L, Cortés-Penagos C, López-Bucio J. *Trichoderma Virens*, A Plant Beneficial Fungus, Enhances biomass Production and Promotes Lateral Root Growth through an Auxin-Dependent Mechanism in *Arabidopsis*. *PlantPhysiol.* 2009; 149(3): 1579–1592.
  - 73. Chowdappa P, Mohan KSP, Jyothi LM, Upreti KK. Growth Stimulation and Induction of Systemic Resistance in Tomato Against Early and Late Blight by *Bacillus Subtilis* OTPB1 or *Trichoderma Harzianum* OTPB3. *Biological Control*. 2013; 65: 109–117.
  - 74. Colla G, Nardi S, Cardarelli M, Ertani A, Lucini L, Canaguierie R, Rouphael Y. Protein Hydrolysates as Biostimulants in Horticulture. *Sci. Hortic.* 2015; 196: 28–38.
  - 75. Velmourougan K, Prasanna R, Chawla G, Nain L, Kumar A, Saxena K. *Trichoderma*–*Azotobacter* Biofilm Inoculation Improves Soil Nutrient Availability and Plant Growth in Wheat and Cotton. *J Basic Microbiol.* 2019; 59: 632–644.
  - 76. Badda N, Yadav K, Kadian N, Aggarwal A. Impact of Arbuscular Mycorrhizal Fungi with *Trichoderma Viride* and *Pseudomonas Fluorescens* on Growth Enhancement of Genetically Modified Bt Cotton (*Bacillus Thuringiensis*). *J Nat*

- Fibers. 2013; 10: 309–325.
- 77. Wilson AR, Nzokou P, Guney D, Kulac S. Growth Response and Nitrogen use Physiology of Fraser Fir (*Abies Fraseri*), Red Pine (*Pinus Resinosa*), and Hybrid Poplar Under Amino Acid Nutrition. *New For.* 2013; 44: 281–295.
  - 78. Colla G, Hoagland L, Ruzzi M, Cardarelli M, Bonini P, Canaguier R, Rouphael Y. Biostimulant Action of Protein Hydrolysates: Unraveling Their Effects on Plant Physiology and Microbiome. *Front. Plant Sci.* 2017; 8:2202.
  - 79. O'Brien JA, Vega A, Bouguyon E, Krouk G, Gojon A, Coruzzi G, Rodrigo AG. Nitrate Transport, Sensing, and Responses in Plants. *Mol Plant.* 2016; 9, 837–856.
  - 80. Miller AJ, Fan X, Shen Q, Smith SJ. Amino Acids and Nitrate as Signals for the Regulation of Nitrogen Acquisition. *J. Exp. Bot.* 2007; 59, 111–119.
  - 81. Fan X, Gordon-Weeks R, Shen QR, Miller AJ. Glutamine Transport and Feedback Regulation of Nitrate Reductase Activity in Barley Roots Leads to Changes in Cytosolic Nitrate Pools. *J. Exp. Bot.* 2006; 57, 1333–1340.
  - 82. Colla G, Svecova E, Rouphael Y, Cardarelli M, Reynaud H, Canaguier R, Planques B. Effectiveness of a Plant-Derived Protein Hydrolysate to Improve Crop Performances under Different Growing Conditions. *Acta Hortic.* 2013; 1009: 175–179.
  - 83. Van Oosten MJ, Pepe O, De Pascale S, Silletti S, Maggio A. The Role of Biostimulants and Bioeffectors as Alleviators of Abiotic Stress in Crop Plants. *Chem. Biol. Technol. Agric.* 2017; 4:5.
  - 84. Santaniello A, Scartazza A, Gresta F, Loreti E, Biasone A, Di Tommaso D, Piaggesi A, Perata P. *Ascophyllum Nodosum* Seaweed Extract Alleviates Drought Stress in *Arabidopsis* by Affecting Photosynthetic Performance and Related Gene Expression. *Front. Plant Sci.* 2017; 8: 1362.
  - 85. Olivares FL, Aguiar NO, Rosa RCC, Canellas LP. Substrate Biofortification in Combination with Foliar Sprays of Plant Growth Promoting Bacteria and Humic Substances Boosts Production of Organic Tomatoes. *Sci. Hortic.* 2015; 183, 100–108.
  - 86. Zancani M, Bertolini A, Petrussa E, Krajňájivá J, Piccolo A. Fulvic Acid Affects Proliferation and Maturation Phases in *Abies Cephalonica* Embryogenic Cells. *J Plant Physiol.* 2011; 168: 1226–1233.
  - 87. Azevedo RA, Lea PJ. Research on Abiotic and Biotic Stress—What Next? *Ann. Appl. Biol.* 2011; 159: 317–319.
  - 88. Bolton MD. Primary Metabolism and Plant Defense-Fuel for the Fire. *Mol Plant Microbe Interact.* 2009; 5, 487-497.
  - 89. Monte E. Understanding *Trichoderma*, between Biotechnology and Microbial Ecology. *Int Microbiol.* 2001;4(1): 1-4.
  - 90. Ertani A, Pizzeghelio D, Altissimo A, Nardi S. Use Of Meat Hydrolyzate Derived From Tanning Residues as Plant Biostimulant for Hydroponically Grown Maize. *J Plant Nutr Soil Sci.* 2013a; 176: 287–296.
  - 91. Apone F, Tito A, Carola A et al. A Mixture of Peptides and Sugars Derived from Plant Cell Walls Increases Plant Defense Responses to Stress and Attenuates Ageing- Associated Molecular Changes in Cultured Skin Cells. *J Biotechnol.* 2010; 145: 367–376.
  - 92. AshrafM, Foolad MR. Roles Of Glycine Betaine and Proline in Improving Plant Abiotic Stress Resistance. *Environ Exp Bot.* 2007; 59: 206–216.

93. Chen THH, Murata N. Glycinebetaine: An Effective Protectant Against Abiotic Stress in Plants. *Trends Plant Sci.* 2008;13: 499–505.
94. Liang XW, Zhang L, Natarajan SK, Beckker DF. Proline Mechanisms of Stress Survival. *Antioxid Redox Sign.* 2013; 19: 998–1011.
95. dos Reis SP, Lima AM, de Souza CRB. Recent Molecular Advances on Downstream Plant Responses to Abiotic Stress. *Int J Mol Sci.* 2012; 13: 8628–8647.
96. Anjum SA, Farooq M, Wang LC et al. Gas Exchange and Chlorophyll Synthesis of Maize Cultivars are Enhanced by Exogenously-Applied Glycinebetaine Under Drought Conditions. *Plant Soil Environ.* 2011a; 57: 326–331.
97. Einset J, Nielson E, Connolly EL et al. Membrane-Trafficking Raba4c Involved in the Effect of Glycine Betaine on Recovery from Chilling Stress in *Arabidopsis*. *Physiol Plant.* 2007; 130: 511–518.
98. Einset J, Winge P, Bones AM, Connolly EL. The FRO2 Ferric Reductase is Required for Glycine Betaine's Effect on Chilling Tolerance in *Arabidopsis* Roots. *Physiol Plant.* 2008; 134: 334–341.
99. Kinnersley AM, Turano FJ. Gamma Aminobutyric Acid (GABA) and Plant Responses to Stress. *Crit Rev Plant Sci.* 2000;19: 479–509.
100. Visconti F, De Paz JM, Bonet L, Jordà M, Quiñones A, Intrigliolo DS. Effects of a Commercial Calcium Protein Hydrolysate on the Salt Tolerance of *Diospyros* Kaki L. Cv. "Rojo Brillante" Grafted on *Diospyros* Lotus L. *Sci Hortic.* 2015;185: 129–38.
101. Zimniewska M, Wladyka-Przybylak M, Mankowski J. Cellulosic Bast Fibers, Their Structure and Properties Suitable for Composite Applications. In: Kalia S, Kaith BS, Kaur I. Editors. Cellulose Fibers: Bio- and Nano-Polymer Composites. *Green Chemistry and Technology* 2011; 97-119.
102. Thyavihalli Girijappa YG, Mavinkere Rangappa S, Parameswaranpillai J, Siengchin S. Natural Fibers as Sustainable and Renewable Resource for Development of Eco-Friendly Composites: A Comprehensive Review. *Front. Mater.* 2019; 6: 226.
103. Silva RA, Santos JL, Oliveira LS, Soares MRS, dos Santos SMS. Biostimulants on Mineral Nutrition and Fiber Quality of Cotton Crop. *R. Bras. Eng. Agríc. Ambiental.* 2016; 20(12): 1062-1066.
104. Belopukhov SL, Daydakova IV, Malinovskaya EA. Research of Effects of Plant Growth Stimulants on Chemical Composition of Long-Fibred Flax at Different Vegetation Stages. *Chemistry and Computational Simulation. Butlerov Communications* 2002; 2(7): 69-72.
105. Marisvelvam R, Athinarayanan G, Ranjitsingh AJA, Usha Raja Nanthini A, Krishnamoorthy R, Alshatwi A.A. Extraction of Dyes from *Petrocarpus santalinus* and Dyeing of Natural Fibres Using Different Mordants. *FIBRES & TEXTILES in Eastern Europe* 2018; 26, 5(131): 20-23. DOI: 10.5604/01.3001.0011.7312.
106. Yavas A, Avinc O, Gedik G. Ultrasound and Microwave Aided Natural Dyeing of Nettle Biofibre (*Urtica dioica* L.) with Madder (*Rubia tinctorum* L.) *FIBRES & TEXTILES in Eastern Europe* 2017; 25, 4(124): 111-120. DOI: 10.5604/01.3001.0010.2856
107. Kopania E, Wietecha J, Ciechańska D. Studies on Isolation of Cellulose Fibres from Waste Plant Biomass. *FIBRES & TEXTILES in Eastern Europe* 2012; 20, 6B (96): 167-172.

108. Karahan M, Ozkan F, Yildirim K, Karahan N. Investigation of the Properties of Natural Fibre Woven Fabrics as a Reinforcement Materials for Green Composites. *FIBRES & TEXTILES in Eastern Europe* 2016; 24, 4(118): 98-104. DOI: 10.5604/12303666.1201138.
109. Thomas S, Paul SA, Pothan LA, Deepa B. Natural Fibres: Structure, Properties and Applications. In: Kalia S, Kaith BS, Kaur I. editors. *Cellulose Fibers: Bio- and Nano-Polymer Composites. Green Chemistry and Technology* 2011; 4-28.
110. Szparaga A, Kocira S, Kocira A, Czerwińska E, Świeca M, Lorencowicz E, Kornas R, Koszel M, Oniszczuk T. Modification of Growth, Yield, and the Nutraceutical and Antioxidative Potential of Soybean Through the use of Synthetic Biostimulants. *Front. Plant Sci.* 2018; 9:1401.
111. Sahari J, Sapuan SM, Ismarrubie ZN, Rahman MZA. Physical and Chemical Properties of Different Morphological Parts of Sugar Palm Fibres. *FIBRES & TEXTILES in Eastern Europe* 2012; 20, 2(91): 21-24.
112. Frydrych I, Raczyńska M, Cekus Z. Measurement of Cotton Fineness and Maturity by Different Methods. *FIBRES & TEXTILES in Eastern Europe* 2010; 18, 6 (83): 54-59.
113. Kocira S, Szparaga A, Kocira A, Czerwińska E, Depo K, Erlichowska B, Deszcz E. Effect of Applying a Biostimulant Containing Seaweed and Amino Acids on the Content of Fiber Fractions in Three Soybean Cultivars. *Legume Res.* 2019; 42(3): 341-347.
114. Nardi S, Pizzeghello D, Schiavon M, Ertani A. Plant Biostimulants: Physiological Responses Induced by Protein Hydrolyzed-Basedproducts and Humic Substances in Plant Metabolism. *Sci. Agric.* 2016; 73(1): 18-23.
115. Wang J, Wang H, Zhao P, Han L, Jiao G, Zheng Y, Huang S, Xia G. Overexpression of a Profilin (GhPFN2) Promotes the Progression of Developmental Phases in Cotton Fibers. *Plant Cell Physiol.* 2010; 51: 1276-1290.
116. Hanafy Ahmed AH, Darwish E, Hamoda SA, AlObaidy MG. Effect of Putrescine and Humic Acid on Growth, Yield and Chemical Composition of Cotton Plants Grown Under Saline Soil Conditions. *Am-Euras. J. Agric. & Environ. Sci.* 2013; 13 (4): 479-497.
117. Bakry BA, Taha MH, Abdelgawad ZA, Abdallah MMS. The Role of Humic Acid and Proline on Growth, Chemical Constituents and Yield Quantity and Quality of Three Flax Cultivars Grown Under Saline Soil Conditions. *Agric Sci.* 2014; 5: 1566-1575.
118. Dienes D, Egyházi A, Réczey K. Treatment of Recycled Fiber with Trichoderma Cellulases. *Ind Crops Prod.* 2004; 20: 11-21.
119. Pere J, Puolakka A, Nousiainen P, Buchert J. Action of Purified *Trichoderma Reesei* Cellulases on Cotton Fibers and Yarn. *J Biotechnol.* 2001; 89(2-3): 247-255.
120. de França Passos D, Pereira Jr. N, Machado de Castro A. A Comparative Review of Recent Advances in Cellulases Production by *Aspergillus*, *Penicillium* and *Trichoderma* Strains and their use for Lignocellulose Deconstruction. *Current Opinion in Green and Sustainable Chemistry* 2018; 14: 60-66.
121. Al-Samaraee WH, Ahmed AA, Hussein HZ, Alwaeli SN. Effect of *Trichoderma Harzianum*, on Chemical Composition and In Vitro Digestibility of Crop Residues. *Plant Archives* 2019; 19(2): 3623-3628.

122. Prasad BMVS, Bhattiprolu SL, Kumari VP, Kumar PA. Study of Antagonistic Capabilities of *Trichoderma* spp. against *Alternaria macrospora* Zimm. Causing Leaf Spot in Cotton. *Int.J.Curr.Microbiol.App.Sci.* 2018; 7(6): 1146-1154.
123. Gallou A, Cranenbrouck S, Declerck S. *Trichoderma Harzianum* Elicits Defence Response Genes in Roots of Potato Plantlets Challenged by *Rhizoctonia Solani*. *Eur J Plant Pathol.* 2009; 124(2): 219-230.
124. Kumar N, Tripathi UK. In Vitro Efficacy of *Trichoderma* Spp. and Plant Extracts on *Alternaria Lini* Cause Blight Disease in Linseed (*Linum Usitatissimum* L.). *J. Pharmacogn. Phytochem.* 2018; 7(2): 1478-1482.
125. Meena B, Ramjegathesh R, Ramyabharathi SA. Evaluation of Biocontrol Agents and Fungicides against Stem Bleeding Disease of Coconut. *J. Plant.Crops.* 2014; 42(3): 395-399.
126. Bhattacharyya SK, Sen K, De RK, Bandopadhyay A, Sengupta C, Adhikary NK. Integration of Biocontrol Agents with Fungicide, Weedicide and Plant Growth Regulator for Management of Stem and Root Rot of Jute. *J. Nat. Appl. Sci.* 2017; 9(2): 899 -904.