

## References

1. Liu L, Cao J, Huang J, Cai Y and Yao J. Extraction of pectins with different degrees of esterification from mulberry branch bark. *Bioresource Technology* 2010; 101, 9: 3268–3273.
2. Du J, He ZD, Jiang RW, Ye WC, Xu HX and But PP. Antiviral flavonoids from the root bark of *Morus alba* L. *Phytochemistry* 2003; 62, 8: 1235–1238.
3. Li RJ, Fei JM, Cai YR, Li YF, Feng JQ and Yao JM. Cellulose whiskers extracted from mulberry: a novel biomass production. *Carbohydrate Polymers* 2009; 76, 1: 94–99.
4. Liu L, Jiang T and Yao J. A two-step chemical process for the extraction of cellulose fiber and pectin from mulberry branch bark efficiently. *Journal of Polymers and the Environment* 2011; 19, 3: 568-573.
5. Jang YS, Amna T, Hassan MS, Kim HC, Kim JH, Baik SH, Khil MS. Nanotitania/mulberry fibers as novel textile with anti-yellowing and intrinsic antimicrobial properties. *Ceramics International* 2015; 41, 5: 6274-6280.
6. Qu C and Wang S. Macro-micro structure, antibacterial activity, and physico-mechanical properties of the mulberry bast fibers. *Fibers and Polymers* 2011; 12, 4: 471-477.
7. Qu C, Wang S, Wang K and Ma Q. Preparation and antibacterial property of the mulberry based textiles. *Fibers and Polymers* 2014; 15, 3: 498-503.
8. Park TY and Lee SG. A study on coarse Hanji yarn manufacturing and properties of the Hanji fabric. *Fibers and Polymers* 2013; 14, 2: 311-315.
9. Wu HL, Li DM, Wu CC, Yang PP and Jing XQ. Research on the structure and properties of mulberry fiber. *Journal of Donghua University* (Eng. Ed.) 2008; 25, 2: 153-158.
10. Brühlmann F, Leupin M, Erismann KH and Fiechter A. Enzymatic degumming of ramie bast fibers. *Journal of Biotechnology* 2000; 76, 1: 43-50.
11. Das B, Chakrabarti K, Ghosh S, Majumdar B, Tripathi S and Chakraborty A. Effect of efficient pectinolytic bacterial isolates on retting and fibre quality of jute. *Industrial Crops and Products* 2012; 36, 1: 415-419.
12. Zheng L, Du Y and Zhang J. Degumming of ramie fibers by alkalophilic bacteria and their polysaccharide-degrading enzymes. *Bioresource Technology* 2001; 78, 1: 89-94.
13. Cao J, Zheng L and Chen S. Screening of pectinase producer from alkalophilic bacteria and study on its potential application in degumming of ramie. *Enzyme and Microbial Technology* 1992; 14, 12: 1013-1016.
14. Kapoor M, Beg QK, Bhushan B, Singh K, Dadhich KS and Hoondal GS. Application of an alkaline and thermostable polygalacturonase from *Bacillus* sp. MG-cp-2 in degumming of ramie (*Boehmeria nivea*) and sunn hemp (*Crotalaria juncea*) bast fibres. *Process Biochemistry* 2001; 36, 8: 803-807.
15. Jacob N, Niladevi KN, Anisha GS and Prema P. Hydrolysis of pectin: an enzymatic approach and its application in banana fiber processing. *Microbiological Research* 2008; 163, 5: 538-544.
16. Kobayashi T, Higaki N, Suzumatsu A, Sawada K, Hagihara H, Kawai S and Ito S. Purification and properties of a high-molecular-weight, alkaline exopolysaccharidase from a strain of *Bacillus*. *Enzyme and Microbial Technology* 2001; 29, 1: 70-75.
17. Bailey MJ, Biely P and Poutanen K. Interlaboratory testing of methods for assay of xylanase activity. *Journal of Biotechnology* 1992; 23, 3: 257–270.

18. Miller GL, Blum R, Glennon WE and Burton AL. Measurement of carboxymethylcellulase activity. *Analytical Biochemistry* 1960; 1, 2: 127-132.
19. Dong Z, Hou X, Sun F, Zhang L and Yang Y. Textile grade long natural cellulose fibers from bark of cotton stalks using steam explosion as a pretreatment. *Cellulose* 2014; 21, 5: 3851-3860.
20. Yim KY, Kan CW. A Comparison Study of Fabric Objective Measurement (FOM) Using KES-FB and PhabrOmeter System on Warp Knitted Fabrics Handle–Smoothness, Stiffness and Softness. World Academy of Science, Engineering and Technology. *International Journal of Chemical, Molecular, Nuclear, Materials and Metallurgical Engineering* 2014; 8, 8: 789-792.
21. Reddy N and Yang Y. Properties and potential applications of natural cellulose fibers from the bark of cotton stalks. *Bioresource Technology* 2009; 100, 14: 3563-3569.
22. French A. Idealized powder diffraction patterns for cellulose polymorphs. *Cellulose* 2014; 21, 2: 885–896.
23. Mora'n JI, Alvarez VA, Cyras VP, Va'zquez A. Extraction of cellulose and preparation of nanocellulose from sisal fibers. *Cellulose* 2008; 15, 1: 149–159.
24. Jayaramudu J, Guduri BR and Rajulu AV. Characterization of new natural cellulosic fabric *Grewia tilifolia*. *Carbohydrate Polymers* 2010; 79, 4: 847-851.